



KING EDWARD VI  
HANDSWORTH GRAMMAR  
SCHOOL FOR BOYS



KING EDWARD VI  
ACADEMY TRUST  
BIRMINGHAM

# Year 12

## Pure Mathematics

### P1 7 Algebraic Methods

## Booklet

HGS Maths



Dr Frost Course



Name: \_\_\_\_\_

Class: \_\_\_\_\_

## Contents

[7.2 Dividing Polynomials](#)

[7.3 The Factor Theorem](#)

**Extract from Formulae booklet**

**Past Paper Practice**

**Summary**

## Prior knowledge check

### Prior knowledge check

**1** Simplify:

**a**  $3x^2 \times 5x^5$       **b**  $\frac{5x^3y^2}{15x^2y^3}$       ← Section 1.1

**2** Factorise:

**a**  $x^2 - 2x - 24$       **b**  $3x^2 - 17x + 20$   
← Section 1.3

**3** Use long division to calculate:

**a**  $197\,041 \div 23$       **b**  $56\,168 \div 34$   
← GCSE Mathematics

**4** Find the equations of the lines that pass through these pairs of points:

**a**  $(-1, 4)$  and  $(5, -14)$   
**b**  $(2, -6)$  and  $(8, -3)$       ← GCSE Mathematics


**5** Complete the square for the expressions:

**a**  $x^2 - 2x - 20$       **b**  $2x^2 + 4x + 15$   
← Section 2.2


PR work from 7.1 on DFM:

Use K201a, K201b, K201c, K201d and K201e

## 7.2 Dividing Polynomials



$$\frac{f(x)}{\text{divisor}(d)} = \text{quotient } (q) + \frac{\text{remainder } (r)}{\text{divisor } (d)}$$



### STEPS:

1. Pick first part of  $q$  by making highest power
2. Fill in rest of column
3. Pick next part of  $q$  to balance next highest power
4. Repeat until  $q$  is a constant term

If you divide cubic  $f(x)$  by  $(ax + b)$  then:

- If  $(ax + b)$  IS A FACTOR then:  $f(x) = (ax + b)(ax^2 + bx + c)$  where  $ax^2 + bx + c$  is the quotient from above
- IF IT IS NOT A FACTOR then the remainder is found by calculating  $f(-\frac{b}{a})$

# Notes

## Exercise 7B

1 Write each polynomial in the form  $(x \pm p)(ax^2 + bx + c)$  by dividing:

a  $x^3 + 6x^2 + 8x + 3$  by  $(x + 1)$

b  $x^3 + 10x^2 + 25x + 4$  by  $(x + 4)$

c  $x^3 - x^2 + x + 14$  by  $(x + 2)$

d  $x^3 + x^2 - 7x - 15$  by  $(x - 3)$

e  $x^3 - 8x^2 + 13x + 10$  by  $(x - 5)$

f  $x^3 - 5x^2 - 6x - 56$  by  $(x - 7)$

2 Write each polynomial in the form  $(x \pm p)(ax^2 + bx + c)$  by dividing:

a  $6x^3 + 27x^2 + 14x + 8$  by  $(x + 4)$

b  $4x^3 + 9x^2 - 3x - 10$  by  $(x + 2)$

c  $2x^3 + 4x^2 - 9x - 9$  by  $(x + 3)$

d  $2x^3 - 15x^2 + 14x + 24$  by  $(x - 6)$

e  $-5x^3 - 27x^2 + 23x + 30$  by  $(x + 6)$

f  $-4x^3 + 9x^2 - 3x + 2$  by  $(x - 2)$

3 Divide:

a  $x^4 + 5x^3 + 2x^2 - 7x + 2$  by  $(x + 2)$

b  $4x^4 + 14x^3 + 3x^2 - 14x - 15$  by  $(x + 3)$

c  $-3x^4 + 9x^3 - 10x^2 + x + 14$  by  $(x - 2)$

d  $-5x^5 + 7x^4 + 2x^3 - 7x^2 + 10x - 7$  by  $(x - 1)$

4 Divide:

a  $3x^4 + 8x^3 - 11x^2 + 2x + 8$  by  $(3x + 2)$

b  $4x^4 - 3x^3 + 11x^2 - x - 1$  by  $(4x + 1)$

c  $4x^4 - 6x^3 + 10x^2 - 11x - 6$  by  $(2x - 3)$

d  $6x^5 + 13x^4 - 4x^3 - 9x^2 + 21x + 18$  by  $(2x + 3)$

e  $6x^5 - 8x^4 + 11x^3 + 9x^2 - 25x + 7$  by  $(3x - 1)$

f  $8x^5 - 26x^4 + 11x^3 + 22x^2 - 40x + 25$  by  $(2x - 5)$

g  $25x^4 + 75x^3 + 6x^2 - 28x - 6$  by  $(5x + 3)$

h  $21x^5 + 29x^4 - 10x^3 + 42x - 12$  by  $(7x - 2)$

5 Divide:

a  $x^3 + x + 10$  by  $(x + 2)$

b  $2x^3 - 17x + 3$  by  $(x + 3)$

c  $-3x^3 + 50x - 8$  by  $(x - 4)$

**Hint** Include  $0x^2$  when you write out  $f(x)$ .

6 Divide:

a  $x^3 + x^2 - 36$  by  $(x - 3)$

b  $2x^3 + 9x^2 + 25$  by  $(x + 5)$

c  $-3x^3 + 11x^2 - 20$  by  $(x - 2)$

## Practice Book

**1** Write each polynomial in the form  $(x \pm p)(ax^2 + bx + c)$  by dividing:

**a**  $2x^3 - 5x^2 + 8x - 5$  by  $(x - 1)$

**b**  $3x^3 + 8x^2 + 3x - 2$  by  $(x + 2)$

**c**  $2x^3 + x^2 - 17x - 12$  by  $(x - 3)$

**d**  $4x^3 + 13x^2 - 11x + 4$  by  $(x + 4)$

**2** Divide:

**a**  $3x^4 + 8x^3 - x^2 - 13x - 6$  by  $(x + 2)$

**b**  $4x^4 - 8x^3 + x^2 - x - 2$  by  $(2x + 1)$

**c**  $9x^4 - 3x^3 - 17x^2 + 13x - 2$  by  $(3x - 2)$

**d**  $4x^4 - 12x^3 - 5x^2 + 15x + 9$  by  $(2x - 3)$

## Worked Example

Find the remainder when  $2x^3 - 5x^2 - 16x + 10$  is divided by  $(x - 4)$ .

## Worked Example

Divide  $8x^3 - 1$  by  $(2x - 1)$ .

## Exercise 7B

7 Show that  $x^3 + 2x^2 - 5x - 10 = (x + 2)(x^2 - 5)$

8 Find the remainder when:

- a  $x^3 + 4x^2 - 3x + 2$  is divided by  $(x + 5)$       b  $3x^3 - 20x^2 + 10x + 5$  is divided by  $(x - 6)$   
 c  $-2x^3 + 3x^2 + 12x + 20$  is divided by  $(x - 4)$

9 Show that when  $3x^3 - 2x^2 + 4$  is divided by  $(x - 1)$  the remainder is 5.

10 Show that when  $3x^4 - 8x^3 + 10x^2 - 3x - 25$  is divided by  $(x + 1)$  the remainder is  $-1$ .

11 Show that  $(x + 4)$  is a factor of  $5x^3 - 73x + 28$ .

12 Simplify  $\frac{3x^3 - 8x - 8}{x - 2}$

**Hint** Divide  $3x^3 - 8x - 8$  by  $(x - 2)$ .

13 Divide  $x^3 - 1$  by  $(x - 1)$ .

**Hint** Write  $x^3 - 1$  as  $x^3 + 0x^2 + 0x - 1$ .

14 Divide  $x^4 - 16$  by  $(x + 2)$ .

**E/P** 15  $f(x) = 10x^3 + 43x^2 - 2x - 10$

Find the remainder when  $f(x)$  is divided by  $(5x + 4)$ .      **(2 marks)**

**E/P** 16  $f(x) = 3x^3 - 14x^2 - 47x - 14$

- a Find the remainder when  $f(x)$  is divided by  $(x - 3)$ .      **(2 marks)**  
 b Given that  $(x + 2)$  is a factor of  $f(x)$ , factorise  $f(x)$  completely.      **(4 marks)**

**Problem-solving**

Write  $f(x)$  in the form  $(x + 2)(ax^2 + bx + c)$  then factorise the quadratic factor.

**E/P** 17 a Find the remainder when  $x^3 + 6x^2 + 5x - 12$  is divided by

- i  $x - 2$ ,  
 ii  $x + 3$ .

**(3 marks)**

b Hence, or otherwise, find all the solutions to the equation  $x^3 + 6x^2 + 5x - 12 = 0$ .      **(4 marks)**

**E/P** 18  $f(x) = 2x^3 + 3x^2 - 8x + 3$

- a Show that  $f(x) = (2x - 1)(ax^2 + bx + c)$  where  $a$ ,  $b$  and  $c$  are constants to be found.      **(2 marks)**  
 b Hence factorise  $f(x)$  completely.      **(4 marks)**  
 c Write down all the real roots of the equation  $f(x) = 0$ .      **(2 marks)**

**E/P** 19  $f(x) = 12x^3 + 5x^2 + 2x - 1$

- a Show that  $(4x - 1)$  is a factor of  $f(x)$  and write  $f(x)$  in the form  $(4x - 1)(ax^2 + bx + c)$ .      **(6 marks)**  
 b Hence, show that the equation  $12x^3 + 5x^2 + 2x - 1 = 0$  has exactly 1 real solution.      **(2 marks)**

# Practice Book

3 Divide:

- a  $2x^3 + 6x^2 - 4$  by  $(x + 1)$
- b  $3x^3 + 7x^2 + 18$  by  $(x + 3)$
- c  $4x^3 - 11x - 10$  by  $(x - 2)$
- d  $2x^3 + 7x^2 + 75$  by  $(x + 5)$

4 Find the remainder when:

- a  $x^3 + 3x^2 + 5x - 8$  is divided by  $(x + 4)$
- b  $2x^3 - 5x^2 + 12x - 20$  is divided by  $(x - 3)$
- c  $3x^3 + 2x^2 - 40x + 45$  is divided by  $(x + 5)$

**E** 5 Find the remainder when  $-15x^3 + 26x^2 - 13x + 5$  is divided by  $(5x - 2)$ . (2 marks)

**E/P** 6  $f(x) = 6x^3 - 13x^2 - 13x + 30$

a Find the remainder when  $f(x)$  is divided by  $(x + 3)$ . (2 marks)

b Given that  $(x - 2)$  is a factor of  $f(x)$ , factorise  $f(x)$  completely. (4 marks)

**E/P** 7  $f(x) = 2x^3 + 3x^2 - 4x + k$  where  $k$  is a constant.

Given that  $(x + 3)$  is a factor of  $f(x)$ :

a find the value of  $k$  (2 marks)

b express  $f(x)$  in the form  $(x + 3)(ax^2 + bx + c)$  where  $a$ ,  $b$  and  $c$  are constants (2 marks)

c show that  $f(x) = 0$  has exactly one real solution. (2 marks)

**E/P** 8  $f(x) = 3x^3 + 10x^2 - 8x - 5$

a Find the remainder,  $r$ , when  $f(x)$  is divided by  $(x - 2)$ . (2 marks)

b Express  $f(x)$  in the form  $(x - 2)(ax^2 + bx + c) + r$  where  $a$ ,  $b$ ,  $c$  and  $r$  are constants. (4 marks)

**E/P** 9  $f(x) = 10x^3 - 29x^2 + 4x + 15$

a Given that  $(x - 1)$  is a factor of  $f(x)$ , express  $f(x)$  in the form  $(x - 1)(ax^2 + bx + c)$ , where  $a$ ,  $b$  and  $c$  are constants. (2 marks)

b Hence factorise  $f(x)$  completely. (4 marks)

c Write down all the solutions to the equation  $f(x) = 0$ . (2 marks)

**Hint** Write the polynomial in part a as  $2x^3 + 6x^2 + 0x - 4$  before dividing.

**Hint** If there is a remainder, then the linear expression  $(x \pm p)$  is not a factor. The polynomial can be written as  $(x \pm p)(ax^2 + bx + c) + r$  where  $r$  is the remainder.

### 7.3 The Factor Theorem

The Factor Theorem states that if  $f(x)$  is a polynomial then:

- If  $f\left(\frac{b}{a}\right) = 0$ , then  $(ax - b)$  is a factor of  $f(x)$ .
- Conversely, if  $(ax - b)$  is a factor of  $f(x)$ , then  $f\left(\frac{b}{a}\right) = 0$ .

- When showing something is a factor in exam questions state 'by factor theorem'

## Notes

## Worked Example

498f: Use the factor theorem to find a unknown coefficient.

$$f(x) = 4x^3 + 12x^2 - 19x + a \text{ where } a \text{ is a constant}$$

Given that  $(2x - 3)$  is a factor of  $f(x)$ , find the value of  $a$ .

## Worked Example

498g: Use the factor theorem to find two unknown coefficients.

Given that  $(x + 4)$  and  $(x + 3)$  are factors of  $f(x) = x^3 + ax^2 - 2x + b$ , determine the values of the constants  $a$  and  $b$ .

$$a = \boxed{\phantom{000}}, \quad b = \boxed{\phantom{000}}$$

## Worked Example

500a: Solve cubic equations using the factor theorem, given one of the roots.

Given that

$$x = -6$$

is a solution to the equation

$$2x^3 - x^2 - 58x + 120 = 0$$

find all the solutions to the equation.

## Worked Example

- a) Fully factorise  $2x^3 + x^2 - 18x - 9$   
b) Hence sketch the graph of  $y = 2x^3 + x^2 - 18x - 9$

## Exam Q

P1 2022

2.  $f(x) = (x - 4)(x^2 - 3x + k) - 42$  where  $k$  is a constant

Given that  $(x + 2)$  is a factor of  $f(x)$ , find the value of  $k$ .

(3)

## Your Turn

2.  $f(x) = (x - 2)(x^3 + 8x^2 - 20x + k) - 1225$  where  $k$  is a constant

Given that  $(x + 3)$  is a factor of  $f(x)$ , find the value of  $k$ .

**(3)**

**(Total for Question 2 is 3 marks)**

## Exam Q

P1 2019

1. 
$$f(x) = 3x^3 + 2ax^2 - 4x + 5a$$

Given that  $(x + 3)$  is a factor of  $f(x)$ , find the value of the constant  $a$ .

**(3)**

## Your Turn

1.

$$f(x) = x^3 + 2bx^2 - 6x - b$$

Given that  $(x + 1)$  is a factor of  $f(x)$ , find the value of the constant  $b$ .

**(Total for Question 1 is 3 marks)**

## Exercise 7C

**1** Use the factor theorem to show that:

- a**  $(x - 1)$  is a factor of  $4x^3 - 3x^2 - 1$       **b**  $(x + 3)$  is a factor of  $5x^4 - 45x^2 - 6x - 18$   
**c**  $(x - 4)$  is a factor of  $-3x^3 + 13x^2 - 6x + 8$ .

**2** Show that  $(x - 1)$  is a factor of  $x^3 + 6x^2 + 5x - 12$  and hence factorise the expression completely.

**3** Show that  $(x + 1)$  is a factor of  $x^3 + 3x^2 - 33x - 35$  and hence factorise the expression completely.

**4** Show that  $(x - 5)$  is a factor of  $x^3 - 7x^2 + 2x + 40$  and hence factorise the expression completely.

**5** Show that  $(x - 2)$  is a factor of  $2x^3 + 3x^2 - 18x + 8$  and hence factorise the expression completely.

**6** Each of these expressions has a factor  $(x \pm p)$ . Find a value of  $p$  and hence factorise the expression completely.

- a**  $x^3 - 10x^2 + 19x + 30$       **b**  $x^3 + x^2 - 4x - 4$       **c**  $x^3 - 4x^2 - 11x + 30$

**7 i** Fully factorise the right-hand side of each equation.

**ii** Sketch the graph of each equation.

- a**  $y = 2x^3 + 5x^2 - 4x - 3$       **b**  $y = 2x^3 - 17x^2 + 38x - 15$       **c**  $y = 3x^3 + 8x^2 + 3x - 2$   
**d**  $y = 6x^3 + 11x^2 - 3x - 2$       **e**  $y = 4x^3 - 12x^2 - 7x + 30$

**(P)** **8** Given that  $(x - 1)$  is a factor of  $5x^3 - 9x^2 + 2x + a$ , find the value of  $a$ .

**(P)** **9** Given that  $(x + 3)$  is a factor of  $6x^3 - bx^2 + 18$ , find the value of  $b$ .

**(P)** **10** Given that  $(x - 1)$  and  $(x + 1)$  are factors of  $px^3 + qx^2 - 3x - 7$ , find the values of  $p$  and  $q$ .

**(P)** **11** Given that  $(x + 1)$  and  $(x - 2)$  are factors of  $cx^3 + dx^2 - 9x - 10$ , find the values of  $c$  and  $d$ .

**(P)** **12** Given that  $(x + 2)$  and  $(x - 3)$  are factors of  $gx^3 + hx^2 - 14x + 24$ , find the values of  $g$  and  $h$ .

**(E)** **13**  $f(x) = 3x^3 - 12x^2 + 6x - 24$

- a** Use the factor theorem to show that  $(x - 4)$  is a factor of  $f(x)$ .      **(2 marks)**  
**b** Hence, show that 4 is the only real root of the equation  $f(x) = 0$ .      **(4 marks)**

**(E)** **14**  $f(x) = 4x^3 + 4x^2 - 11x - 6$

- a** Use the factor theorem to show that  $(x + 2)$  is a factor of  $f(x)$ .      **(2 marks)**  
**b** Factorise  $f(x)$  completely.      **(4 marks)**  
**c** Write down all the solutions of the equation  $4x^3 + 4x^2 - 11x - 6 = 0$ .      **(1 mark)**

**(E)** **15 a** Show that  $(x - 2)$  is a factor of  $9x^4 - 18x^3 - x^2 + 2x$ .      **(2 marks)**

**b** Hence, find four real solutions to the equation  $9x^4 - 18x^3 - x^2 + 2x = 0$ .      **(5 marks)**

### Challenge

$$f(x) = 2x^4 - 5x^3 - 42x^2 - 9x + 54$$

- a** Show that  $f(1) = 0$  and  $f(-3) = 0$ .  
**b** Hence, solve  $f(x) = 0$ .

### Problem-solving

Use the factor theorem to form simultaneous equations.

# Practice Book

1 Use the factor theorem to show that:

- a  $(x + 1)$  is a factor of  $2x^3 + 7x^2 - 5$
- b  $(x + 2)$  is a factor of  $x^3 + 4x^2 + 3x - 2$
- c  $(x - 3)$  is a factor of  $2x^3 - 3x^2 - 7x - 6$
- d  $(x - 4)$  is a factor of  $x^4 - 3x^3 - 15x - 4$

2 Use the factor theorem to show that the linear expression is a factor of the polynomial  $f(x)$  and factorise  $f(x)$  completely:

- a  $(x - 2)$ ,  $2x^3 + x^2 - 13x + 6$
- b  $(x + 3)$ ,  $2x^3 + 17x^2 + 38x + 15$
- c  $(x - 1)$ ,  $6x^3 - x^2 - 11x + 6$
- d  $(x + 4)$ ,  $15x^3 + 61x^2 - 2x - 24$

3 Fully factorise each expression:

- a  $x^3 + 2x^2 - 21x + 18$
- b  $2x^3 + 13x^2 + 13x - 10$
- c  $3x^3 + 2x^2 - 41x - 60$

4 For each of the following polynomials,

- i fully factorise each polynomial  $f(x)$ .
- ii Hence sketch the graph of  $y = f(x)$ .

- a  $2x^3 - 11x^2 + 5x + 18$
- b  $2x^3 - 3x^2 - 39x + 20$
- c  $6x^3 + 37x^2 + 50x - 21$

**Hint** The **factor theorem** states that if

$f(x)$  is a polynomial, then:

- if  $f(p) = 0$  then  $(x - p)$  is a factor of  $f(x)$
- if  $(x - p)$  is a factor of  $f(x)$  then  $f(p) = 0$

**Hint**

When you have used the factor theorem to show the linear expression is a factor, you can use long division to find the quadratic factor. Factorise the quadratic factor to write the polynomial as a product of three linear factors.

**Hint**

Try values of  $p$  in each expression for  $f(x)$ , e.g.  $p = -1, 1, 2, 3, \dots$  until you find  $f(p) = 0$ . Then use the factor theorem to deduce that  $(x - p)$  is a factor of  $f(x)$ .

**Hint**

To sketch the graph, you need to identify the points where the curve crosses the axes. Set  $x = 0$  to find the  $y$ -intercept and  $y = 0$  to find the  $x$ -intercepts.

The general shapes of cubic graphs are:



if the coefficient of  $x^3$  is positive  
if the coefficient of  $x^3$  is negative

← Section 4.1

**E/P 5**  $f(x) = 6x^3 - 17x^2 - 15x + 36$

Given that  $(x - 3)$  is a factor of  $f(x)$ , find all the solutions to  $f(x) = 0$ . **(5 marks)**

**E/P 6**  $f(x) = 9x^3 + 24x^2 - 44x + 16$

a Use the factor theorem to show that  $(x + 4)$  is a factor of  $f(x)$ . **(2 marks)**

b Hence show that  $f(x)$  can be written in the form  $f(x) = (x + 4)(px + q)^2$ , where  $p$  and  $q$  are integers to be found. **(4 marks)**

**E 7**  $f(x) = 2x^3 - 3x^2 - 5x + 6$ . Factorise  $f(x)$  completely. **(5 marks)**

**E/P 8**  $g(x) = x^3 + 2x^2 - 19x + k$

Given that  $(x + 1)$  is a factor of  $g(x)$ ,

- a show that  $k = -20$  **(2 marks)**
- b express  $g(x)$  as a product of three linear factors. **(3 marks)**
- c Sketch the curve with equation  $y = x^3 + 2x^2 - 19x - 20$ , indicating the values where the curve crosses the  $x$ -axis and the  $y$ -axis. **(4 marks)**

**E/P 9**  $p(x) = 25x^3 + 55x^2 - 56x + 12$

- a Use the factor theorem to show that  $(x + 3)$  is a factor of  $p(x)$ . **(2 marks)**
- b Fully factorise  $p(x)$ . **(3 marks)**
- c Hence show that there are exactly two real roots of the equation  $p(x) = 0$ . **(2 marks)**

# Past Paper Questions

## A2 2021 Paper 1

### Algebraic Methods

1.

$$f(x) = ax^3 + 10x^2 - 3ax - 4$$

Given that  $(x - 1)$  is a factor of  $f(x)$ , find the value of the constant  $a$ .

You must make your method clear.

(3)



### Exams

- Formula Booklet
- Past Papers
- Practice Papers
- [past paper Qs by topic](#)

Past paper practice by topic. Both new and old specification can be found via this link on [hgsmaths.com](http://hgsmaths.com)

A2 2021 Paper 1

Algebraic Methods

Question	Scheme	Marks	AOs
1	$f(x) = a(x^3 + 10x^2 - 3ax - 4) = 0$	3.1a	M1
	$6 - 2a = 0 \Rightarrow a = 3$	1.1b	M1
		1.1b	A1
		(3)	

Notes (3 marks)

Main method seen:  
 M1: Attempts  $f(1) = 0$  to set up an equation in  $a$ . It is implied by  $a + 10 - 3a - 4 = 0$   
 Condense a slip but attempting  $f(-1) = 0$  is M0  
 M1: Solves a linear equation in  $a$ .  
 Using the main method it is dependent upon having set  $f(1) = 0$   
 It is implied by a solution of  $\pm a \pm 10 \pm 3a \pm 4 = 0$ .  
 Don't be concerned about the mechanics of the solution.  
 A1:  $a = 3$  (following correct work)

## Summary of Key Points

- 1** When simplifying an algebraic fraction, factorise the numerator and denominator where possible and then cancel common factors.
- 2** You can use long division to divide a polynomial by  $(x \pm p)$ , where  $p$  is a constant.
- 3** The **factor theorem** states that if  $f(x)$  is a polynomial then:
  - If  $f(p) = 0$ , then  $(x - p)$  is a factor of  $f(x)$
  - If  $(x - p)$  is a factor of  $f(x)$ , then  $f(p) = 0$

## Mixed Exercise

1 Simplify these fractions as far as possible:

a  $\frac{3x^4 - 21x}{3x}$

b  $\frac{x^2 - 2x - 24}{x^2 - 7x + 6}$

c  $\frac{2x^2 + 7x - 4}{2x^2 + 9x + 4}$

2 Divide  $3x^3 + 12x^2 + 5x + 20$  by  $(x + 4)$ .

3 Simplify  $\frac{2x^3 + 3x + 5}{x + 1}$

**(E)** 4 a Show that  $(x - 3)$  is a factor of  $2x^3 - 2x^2 - 17x + 15$ . (2 marks)

b Hence express  $2x^3 - 2x^2 - 17x + 15$  in the form  $(x - 3)(Ax^2 + Bx + C)$ , where the values  $A$ ,  $B$  and  $C$  are to be found. (3 marks)

**(E)** 5 a Show that  $(x - 2)$  is a factor of  $x^3 + 4x^2 - 3x - 18$ . (2 marks)

b Hence express  $x^3 + 4x^2 - 3x - 18$  in the form  $(x - 2)(px + q)^2$ , where the values  $p$  and  $q$  are to be found. (4 marks)

**(E)** 6 Factorise completely  $2x^3 + 3x^2 - 18x + 8$ . (6 marks)

**(E/P)** 7 Find the value of  $k$  if  $(x - 2)$  is a factor of  $x^3 - 3x^2 + kx - 10$ . (4 marks)

**(E/P)** 8  $f(x) = 2x^2 + px + q$ . Given that  $f(-3) = 0$ , and  $f(4) = 21$ :

a find the value of  $p$  and  $q$

b factorise  $f(x)$ . (6 marks)

(3 marks)

**(E/P)** 9  $h(x) = x^3 + 4x^2 + rx + s$ . Given  $h(-1) = 0$ , and  $h(2) = 30$ :

a find the values of  $r$  and  $s$

b factorise  $h(x)$ . (6 marks)

(3 marks)

**(E)** 10  $g(x) = 2x^3 + 9x^2 - 6x - 5$ .

a Factorise  $g(x)$ . (6 marks)

b Solve  $g(x) = 0$ . (2 marks)

## Mixed Exercise

- (E) 11 a** Show that  $(x - 2)$  is a factor of  $f(x) = x^3 + x^2 - 5x - 2$ . **(2 marks)**
- b** Hence, or otherwise, find the exact solutions of the equation  $f(x) = 0$ . **(4 marks)**
- (E) 12** Given that  $-1$  is a root of the equation  $2x^3 - 5x^2 - 4x + 3$ , find the two positive roots. **(4 marks)**
- (E) 13**  $f(x) = x^3 - 2x^2 - 19x + 20$
- a** Show that  $(x + 4)$  is a factor of  $f(x)$ . **(3 marks)**
- b** Hence, or otherwise, find all the solutions to the equation  $x^3 - 2x^2 - 19x + 20 = 0$ . **(4 marks)**
- (E) 14**  $f(x) = 6x^3 + 17x^2 - 5x - 6$
- a** Show that  $f(x) = (3x - 2)(ax^2 + bx + c)$ , where  $a$ ,  $b$  and  $c$  are constants to be found. **(2 marks)**
- b** Hence factorise  $f(x)$  completely. **(4 marks)**
- c** Write down all the real roots of the equation  $f(x) = 0$ . **(2 marks)**

## Problem Solving Set B

### Bronze

$f(x) = x^3 - x^2 + px + q$  where  $p$  and  $q$  are integers.

Given that  $(x + 1)$  is a factor of  $f(x)$ ,

**a** show that  $q - p = 2$ . (3 marks)

Given that  $(x + 3)$  is also a factor of  $f(x)$ ,

**b** show that  $q - 3p = 36$ . (3 marks)

**c** Hence find the value of  $p$  and the corresponding value of  $q$ . (2 marks)

**d** Factorise  $f(x)$  completely. (2 marks)

### Silver

$f(x) = 2x^3 - x^2 + px + q$  where  $p$  and  $q$  are integers.

Given that  $(x + 2)$  is a factor of  $f(x)$ ,

**a** show that  $q - 2p - 20 = 0$ . (3 marks)

Given that  $(x - 3)$  is also a factor of  $f(x)$ ,

**b** find the value of  $p$  and the corresponding value of  $q$ . (5 marks)

**c** Factorise  $f(x)$  completely. (2 marks)

### Gold

$f(x) = x^3 + (p + 4)x^2 + 8x + q$  where  $p$  and  $q$  are integers.

Given that  $(x - 2)$  is a factor of  $f(x)$ ,

**a** show that  $4p + q + 40 = 0$ . (3 marks)

Given that  $(x + p)$  is also a factor of  $f(x)$ , and that  $p > 0$ ,

**b** show that  $4p^2 - 8p + q = 0$ . (3 marks)

**c** Hence find the value of  $p$  and the corresponding value of  $q$ . (5 marks)

**d** Factorise  $f(x)$  completely. (2 marks)